

- controlling absence of tannins in faba bean, *Aust J of Agric. Res.*, 59 pp62- 68.
- Peng,S.; F. V. Garcia, R.C. Laza and Cassman.1993.Adjustment for specific leaf wheat improves chlorophyll meters estimate of rice leaf nitrogen concentration. *J. Agron.*, 85 P P900-987.
- Salih., Z.K.N. S. Ghanim and T.A. Abraham.2010. Response two cultivar of peas (*Pisium sativum* L.) for spraying by nutrient solution Marvel. *J. Tikrit.Univ.Coll.of Agric Sci.*, 10 (2) pp129–137.
- Steel., R. G. D. and J.H.Torrie.1960.Principles and Procedures of Statistics.2nded McGraw. Hill book Company., New York. pp481.
- Tetio., F.K.and F.P.Gardner.1988. Responses of maize to plant population density. I.Canopy development, Light and light interception and vegetative growth.*Agron. J.*, 80 pp930-935.
- Thalooth, A.T.,M.M. Tawfik and H. Mohamed .2006.A comparative study on the effect of foliar application of zinc, potassium and magnesium on growth, yield and some chemical constituents of mung bean plant grown under water stress conditions. *World J .Agric. Sic.*, (1) PP 37-46
- Wasfi., Z. 2003. Cultivation of field crops. Aladdin foundation for printing and publishing. Republic Musrarabn. pp2-30.

## **THE EFFECT OF SPRAYING (MS) NUTRIENT SOLUTION AND BORON ON GROWTH, YIELD AND QUALITY OF (*Vicia faba* L.)**

**MAKKIYAH KADHIM ALAG\***

\* Dept. of Field Crop Sci.- Coll. of Agric- Univ. of Baghdad. makaya201050@yahoo.com

### **ABSTRACT**

A field experiment was conducted at farm of Field Crops Depart, College of Agarc, Univ of Baghdad during the seasons 2010/2011 and 2011/2012 to study the effect of spraying (MS) nutrition solution and boron only or together during the important stages of the plant faba beans the growth, yield and quality.

A split plot on in with R. C. B. D. distribution was three replications. Levels of boron used 0,200, 400 mg.L<sup>-1</sup>.they were assigned to the main plots. While the number of spraying in three intervals (spray the plants with water only) and the treatment of two spray times before flowering and at flowering 25% and the treatment three spray times before and at flowering 25% and flowering 50% were assigned as sub plots.

The results showed a significant effect of the number of times spraying (MS) nutrition solution on plant height, biological yield, chlorophyll content, fertility rate, number of pods per plant, number of seeds per pod, seed weight, seed yield, protein and carbohydrates content, except Width pod in both seasons. The

treatment that was sprayed two time give the highest plant 52.82 و 63.33cm, number of branches. plant<sup>-1</sup> 5.03 و 9.00 branch plant, biological yield 10.77 و 14.16 t.ha<sup>-1</sup>, chlorophyll content of 53.23 and 56.26 (Mu.gm.cm<sup>2</sup>)<sup>-1</sup>-number of pods per plant 6.06 و 12.00 Pod ،seed weight 1312 و 1299mg, total seed yield 6.56 and 6.97t.ha<sup>-1</sup>, ratio of protein 26.26 و 25.84% and carbohydrate 24.25 و 25.01% for two seasons respectively. While the treatment that was sprayed three times gave the longest pods 18.19 و 16.22cm ،fertility rate 8.12 و 13.33%، number of seeds.Pod<sup>-1</sup>،4.57 و 3.78 for two seasons respectively, harvest index 63.12% for the first season only. Significantly influenced levels of boron and interaction in most traits, except width Pod in both seasons.

The results showed that the highest seed yield 7.37 و 8.80 t.ha<sup>-1</sup> of significant interaction between two sprays times of (MS) nutrition solution with concentration of 200 mg.L<sup>-1</sup> of the boron during two seasons respectively.

**Key words:** (MS) Nutrient solution ،Boron،Faba bean.، yield، quality.

**Diyala Agricultural Sciences Journal, 7 ( 1 ):121-132.( 2015 ). ISRA impact factor 4.758.**

<http://www.agriculmag.uodiyala.edu.iq>

<http://www.iasj.net/iasj?func=issueTOC&isId=4427&uiLanguage=en>