## ENVIRONMENTAL SENSITIVITY OF DESERT SOILTO EROSION IN SOUTHERN AL-QADISIYAH LAKE USING GIS.

## Salah M. F. Al-Juraisy University of Al-Anbar/ College of Agriculture/ Soil Science Dept. salahmrsh@yahoo.com

## ABSTRACT

Characterization of soil Sensitivity to erosion in desert ecosystem is very important to conserving fragile ecosystems and planning management and protect ecosystem from degradation . desert ecosystem degradation terms refer to soil erosion, mass movement and basin sedimentation.

The study area located in the southern part of Al- Qadisiyah lake (109.15 ha) at latitude from  $34^{\circ}00N$  to  $34^{\circ}30$  N and longitude from  $42^{\circ}00E$  to  $42^{\circ}20$  E.

A number of 10 soil site were studied. The soil laboratory analyses were: soil texture, Calcium carbonate content, and average of organic matter content was 0.5%) .The results showed That area of study were classified into four main site varied to soil Sensitivity to erosion which are classified to: law sensitive (12.13%), moderate sensitive (26.90%), high sensitive (39.76%),and very high sensitive (21.21%)

Key words: Soil erosion, Environmental Sensitivity, Desert soil, Al-Qadisiyah lake.